

Original Paper

Heterogeneity of kinase inhibitor resistance mechanisms in GIST

B Liegl,^{1,2} I Kepten,³ C Le,³ M Zhu,¹ GD Demetri,⁴ MC Heinrich,^{5,6} CDM Fletcher,¹ CL Corless³ and JA Fletcher^{1*}

¹Department of Pathology, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, USA

²Department of Pathology, Medical University of Graz, Graz, Austria

³Department of Pathology, Oregon Health and Science University, Portland, OR, USA

⁴Ludwig Center, Dana-Farber Cancer Institute and Harvard Medical School, Boston, MA, USA

⁵Division of Hematology and Oncology, Oregon Health and Science University Cancer Institute, Portland, OR, USA

⁶Portland VA Medical Center, Portland, OR, USA

*Correspondence to:

JA Fletcher, Department of Pathology, Brigham and Women's Hospital, 75 Francis Street, Boston, MA 02115, USA.
E-mail: jfletcher@partners.org or to B Liegl-Atzwanger, Department of Pathology, Medical University of Graz, Auenbruggerplatz 25, A-8036 Graz, Austria.

E-mail: bernadette.liegl-atzwanger@meduni-graz.at

No conflicts of interest were declared.

Abstract

Most GIST patients develop clinical resistance to KIT/PDGFR tyrosine kinase inhibitors (TKI). However, it is unclear whether clinical resistance results from single or multiple molecular mechanisms in each patient. KIT and PDGFRA mutations were evaluated in 53 GIST metastases obtained from 14 patients who underwent surgical debulking after progression on imatinib or sunitinib. To interrogate possible resistance mechanisms across a broad biological spectrum of GISTs, inter- and intra-lesional heterogeneity of molecular drug-resistance mechanisms were evaluated in the following: conventional KIT (CD117)-positive GISTs with KIT mutations in exon 9, 11 or 13; KIT-negative GISTs; GISTs with unusual morphology; and KIT/PDGFR wild-type GISTs. Genomic KIT and PDGFRA mutations were characterized systematically, using complementary techniques including D-HPLC for KIT exons 9, 11–18 and PDGFRA exons 12, 14, 18, and mutation-specific PCR (V654A, D820G, N822K, Y823D). Primary KIT oncogenic mutations were found in 11/14 patients (79%). Of these, 9/11 (83%), had secondary drug-resistant KIT mutations, including six (67%) with two to five different secondary mutations in separate metastases, and three (34%) with two secondary KIT mutations in the same metastasis. The secondary mutations clustered in the KIT ATP binding pocket and kinase catalytic regions. FISH analyses revealed KIT amplicons in 2/10 metastases lacking secondary KIT mutations. This study demonstrates extensive intra- and inter-lesional heterogeneity of resistance mutations and gene amplification in patients with clinically progressing GIST. KIT kinase resistance mutations were not found in KIT/PDGFR wild-type GISTs or in KIT-mutant GISTs showing unusual morphology and/or loss of KIT expression by IHC, indicating that resistance mechanisms are fundamentally different in these tumours. Our observations underscore the heterogeneity of clinical TKI resistance, and highlight the therapeutic challenges involved in salvaging patients after clinical progression on TKI monotherapies.

Copyright © 2008 Pathological Society of Great Britain and Ireland. Published by John Wiley & Sons, Ltd.

Keywords: imatinib; sunitinib; drug resistance mechanisms; GIST; heterogeneity

Received: 26 February 2008

Revised: 25 April 2008

Accepted: 1 May 2008

Introduction

Gastrointestinal stromal tumours (GISTs) are the most common mesenchymal tumours of the gastrointestinal (GI) tract and are refractory to radiation and conventional chemotherapy. 85–90% of GISTs have activating mutations of the *KIT* or *PDGFRA* receptor tyrosine kinase genes [1–3], resulting in oncoproteins that are crucial diagnostic and therapeutic targets in GIST [2,3]. Indeed, therapeutic inhibition of *KIT/PDGFR* kinase activity by imatinib has emerged as the first-line treatment option in patients with

inoperable GIST [4–6]. Notably, imatinib response depends on *KIT/PDGFR* mutational status [7,8]. Patients whose GISTs have *KIT* exon 11 mutations have a higher response rate and longer median survival compared to those with wild-type *KIT/PDGFR* or with *KIT* exon 9 mutations [1]. Complete responses to imatinib in metastatic GIST are rare ($\leq 5\%$) and most responding patients develop secondary resistance [6].

The most common secondary resistance mechanism appears to be mutation of the *KIT* kinase domain; however, additional resistance mechanisms include *KIT/PDGFR* genomic amplifications and activation

of alternative oncogenes [9,10]. Therapeutic options for patients whose GISTs progress on imatinib include dose escalation or treatment with sunitinib malate (SUTENT Pfizer, New York, USA), a Federal Drug Administration (FDA)-approved drug with demonstrated efficacy, acceptable tolerability and safety in a double-blind placebo-controlled Phase III trial [11]. Previous studies have focused on individual imatinib resistance mechanisms in GIST lesions progressing on imatinib therapy, but the heterogeneity of these mutations, in a given patient, remains unclear. Therefore, the aim of this study was to characterize intra- and inter-lesional drug resistance mechanisms in GIST tumour samples obtained during debulking procedures performed on patients with imatinib or sunitinib resistance.

Material and methods

Tumour selection

Fifty-three GIST metastases from 14 patients (12 male and 2 female, age range 50–75 years, median age 62 years) were studied. All patients progressed clinically on imatinib or sunitinib, according to the conventional Southwest Oncology Group/Response Evaluation Criteria in Solid Tumours [12]. All patients underwent resection during 2001–2004 at the Brigham and Women's Hospital, Boston, MA, USA. Imatinib or sunitinib was discontinued within 1 week prior to debulking surgeries. Surgery was performed in five patients progressing after imatinib alone and in nine patients progressing after imatinib and sunitinib treatment. To interrogate possible resistance mechanisms in a broad spectrum of GISTs, we examined not only tumours with typical morphology, but tumours that were KIT (CD117) negative, had unusual morphology or were *KIT/PDGFR*A wild-type. This study was approved by the Institutional Review Board of Brigham and Women's Hospital.

Haematoxylin and eosin-stained sections from 276 paraffin blocks were reviewed to confirm the diagnoses prior to inclusion in the study. Tumour regions from different metastases or different areas within metastases were selected from each patient, with an emphasis on variation in tumour cytology, KIT expression (KIT-positive or KIT-negative) and mitotic activity. In total, 57 tumour areas from 53 metastases were selected. The morphological appearance (spindle cell, epithelioid cell, mixed cell type, unusual morphology), tumour size, location, treatment effects (necrosis, hyalinosis, pseudo-chondroid changes, haemorrhage) were evaluated, as well as mitotic rate [expressed as the number of mitotic figures per 50 high power fields (HPFs) in the most mitotic area, using a $\times 40$ objective and a $\times 10$ ocular, field size 0.25 mm^2] (Table 1). Histological treatment response was scored in each metastasis, using a previously proposed grading scheme [8]: 1, minimal (0–10% response); 2, low (>10% and

<50% responses); 3, moderate (>50% and <90% response); and 4, high (>90% response).

Immunohistochemistry

Immunohistochemical studies for KIT (CD117) (Dako, Carpinteria, CA, USA; polyclonal A4502, 1:250) were performed in all cases without epitope retrieval, as previously described [13]. In cases lacking KIT expression, additional immunohistochemical stains using antibodies against SMA (Sigma, St. Louis, MO, USA; 1A4, 1:20,000); Desmin (Dako; D33, 1:500); Caldesmon (Dako; h-CD, 1:300; heat-induced epitope retrieval) and MYF4 (Novocastra, Burlingame, CA, USA; LO26, 1:600, heat-induced epitope retrieval) were performed. The Envision Plus detection system (Dako) was used for all antibodies. Appropriate positive and negative controls were included.

DNA Extraction and initial mutation screening

The 57 tumour areas of interest were marked and collected from unstained sections by manual tumour tissue dissection. Tumour tissue was deparaffinized as previously reported [14]. Mutational analyses were performed on the extracted genomic DNA, using a combination of polymerase chain reaction (PCR) amplification, denaturing high-performance liquid chromatography (D-HPLC) screening and automated sequencing, as described previously [1,2,15]. *KIT* exons 9, 11, 12, 13, 14, 15, 16, 17, 18 and *PDGFRA* exons 12, 14, 18 were evaluated.

Mutation screening by allele-specific PCR

Five known hotspots for *KIT* secondary resistance mutations [V654A, D820G, N822K (T \rightarrow A and T \rightarrow G), Y823D] were screened by novel allele-specific PCR assays. Details on the development of these assays will be the subject of another report (Kepten *et al*, manuscript in preparation). In brief, mutation-specific forward primers were designed such that the nucleotide substitution of interest was matched by a locked nucleic acid at the 3' end. Amplicons were detected by hydrolysable, dual-labelled probes to exon 13 or exon 17, depending on the site of mutation. PCR conditions were established such that dilutions of GIST DNA samples with known mutations (estimated to be 50% by direct sequencing) were routinely positive down to the level of 1:100 (estimated 0.5% mutant allele). DNA from formalin-fixed, paraffin-embedded normal tissue was either negative or had $C_t > 5$ cycles beyond that of the 1:100 dilution. Dilution controls and normal DNA controls were included in each assay. All abnormal allele-specific assays were repeated at least once.

Fluorescence *in situ* hybridization (FISH)

FISH analyses of *KIT* copy number were performed on $4 \mu\text{m}$ tissue sections that were pre-baked for 2 h at

Table 1. Summary of clinicopathological and molecular findings in 14 patients undergoing debulking procedures for progressing GIST during TKI treatment

Patient no.	Treatment	Age/sex	Location	Size (cm)	Morphology	Mitoses/50HPF	Histological treatment		Primary KIT mutation	Secondary KIT mutation	
							response	IHC		KIT/ IHC	DHPLC
1	IM	57M	Serosal nodule (1)	1.8	Spindle	20	+	+	Exon 9 (insertion AY502-503)	NMF	NMF
			Serosal nodule (2)	2.7	Spindle (perivascular growth)	7	+				
			Serosal nodule (3)	4.0	Spindle (perivascular growth)	18	+				
2	IM, SU	66M	Serosal nodule right lower quadrant	1.7	Mixed (clear CP)	4	+	+	Exon 9 (insertion AY502-503)	NMF	NMF
			Omental nodule	0.7	Spindle	10	+				
			Sigmoid mass (1)	5.0	Epithelioid	3	+				
			Sigmoid mass (2)	5.0	Epithelioid	6	-				
			Periumbilical tumour	12.0	Mixed	9	+				
			Serosal nodule right upper quadrant (1)	2.0	Epithelioid	34	+				
Serosal nodule right upper quadrant (2)	4.0	Epithelioid	33	+							
3	IM, SU	62M	Stomach (1)	24.0	Spindle (palisading)	100	+	+	Exon 11 (deletion EVQWKV554-559)	Exon 17	D816H
			Stomach (2)	24.0	Spindle (palisading)	40	+				
4	IM	50M	Liver metastasis	9.5	Mixed	19	+	+	Exon 11 (deletion KPMYEQWK550-558)	Exon 18	A829P
			Stomach (1)	12.3	Mixed	54	+				
			Stomach (2)	12.3	Epithelioid (giant cells)	26	+				
5	IM, SU	59F	Liver metastasis	6.5	Spindle	7	+	+	Exon 11 (deletion PYD577-579)	Exon 13	V654A
			Mesenteric nodule (1)	0.9	Spindle	22	+				
			Mesenteric nodule (2)	2.9	Spindle	34	+				
			Nodule attached to gallbladder	3.0	Spindle (giant cells)	20	+				
6	IM	75M	Intra-abdominal 1	6.0	Spindle	36	+	+	Exon 11 (deletion VEEINGNNYYIDPTQL560-576)	Exon 13	V654A
			Intra-abdominal 2	5.0	Spindle	88	+				
			Intra-abdominal 2	5.0	Spindle	88	+				
7	IM, SU	67F	Stomach	35.0	Epithelioid (intracytoplasmic inclusions)	2	+	-	Exon 11 (deletion WKV557-559(F))	NMF	NMF
			Liver metastasis	2.7	Epithelioid (intracytoplasmic inclusions and atypia)	13	2				
			Serosal nodule	3.0	Epithelioid (intracytoplasmic inclusions and atypia)	14	2				
8	IM	54M	Abdominal tumour (1)	3.0	Spindle	1	+	+	Exon 11 (deletion WKV557-559C)	NMF	NMF
			Tumour Stomach wall	15.4	Mixed	26	+				
			Abdominal tumour (2)	2.5	Epithelioid	8	+				
			Abdominal tumour (3)	2.5	Epithelioid	8	+				

Table 1. Continued

Patient no.	Treatment	Age/sex	Location	Size (cm)	Morphology	Mitoses/50HPF	Histological treatment		Primary KIT mutation	Secondary KIT mutation	
							response	IHC		KIT mutation	DHPLC
9	IM, SU	75M	Mesenteric nodule 1	1.9	Spindle	16	2	+		Exon 13	V654A
			Skin metastasis	3.5	Spindle	57	1	+		Exon 13	V654A
			Mesenteric nodule 2	3.3	Spindle	72	2	+		Exon 11 (point mutation L576P)	V654A
			Mesenteric nodule 4	1.8	Spindle	24	1	+		Exon 13	V654A
			Mesenteric nodule 3	2.9	Spindle	44	1	+		Exon 13 and 17	V654A, D820G
10	IM	74M	Stomach wall (1)	13.0	Epithelioid (intracytoplasmic inclusions)	15	1	+		NMF	NMF
			Stomach wall (2)		Epithelioid	52	1	+	Exon 13 (point mutation K642E)	NMF	NMF
			Stomach wall (3)		Epithelioid	61	2	+		NMF	NMF
11	IM, SU	63M	Tumour mid-abdomen	3.8	Mixed	1	2	-		NMF	NMF
			Serosal nodule left upper quadrant (1)	3.6	Spindle	1	4	+		Exon 14	T670I
			Serosal nodule ileum (1)	0.5	Spindle	7	1	+		Exon 13 (point mutation K642E)	D816H
			Serosal nodule ileum (2)	1.0	Epithelioid	34	1	+		Exon 17	D816H
			Serosal nodule terminal ileum	0.7	Spindle	1	1	+		Exon 17	D816H
			Serosal nodule left upper quadrant (2)	2.8	Mixed	3	2	+		Exon 17	D816H
12	IM, SU	60M	Omental nodule	4.5	Epithelioid	36	1	+			
			Liver metastasis (1)	2.8	Epithelioid (huge eosinophilic CP)	18	1	+			
			Tumour sigmoid wall	14.0	Epithelioid	15	1	+	NMF		NMF
			Liver metastasis (2)	2.1	Epithelioid (huge eosinophilic CP)	16	1	+			
			Serosal nodule adjacent to bowel	3.5	Spindle (storiform)	1	2	+			
13, NF Type I	IM, SU	44M	Nodule lig. of Tritz	3.0	Spindle	1	3	+	NMF		NMF
			Nodule duodenum	2.5	Spindle (storiform)	2	2	+			
			Mesenteric nodule	15.0	Spindle (storiform)	6	2	+			
			Perigastric mass (1)	1.5	Spindle with atypia	48	1	-			
			Perigastric mass (2)	2.5	Epithelioid	4	1	+			
14	IM, SU	65M	Perigastric mass (3)	2.4	Epithelioid	14	2	+			
			Liver metastasis	19.8	Epithelioid	58	1	+	NMF		NMF
			Large nodule lesser curvature (1)	3.5	Epithelioid with mild atypia	19	1	+			
			Large nodule lesser curvature (2)	3.5	Spindle with atypia	60	1	-			

pt, patient; IM, imatinib; SU, sunitinib; CP, cytoplasm; +, positive; -, negative; NMF, no mutation found.

All listed samples were examined for KIT mutations in exons 9, 11, 12, 13, 14, 15, 16, 17, 18 and PDGFRA mutations in exons 12, 14, 18. Exons not specifically listed are wild-type.

Heterogeneity of resistance mechanisms in GIST

60°C. The sections were deparaffinized in xylene three times (each 15 min) and dehydrated twice in 100% ethanol for 2 min. The slides were then immersed in TRIS-EDTA (100 mM Tris base, 50 mM EDTA, pH 7.0) for 45 min at 95–99°C and rinsed in \times 1 PBS for 5 min. Proteolytic digestion of the sections was performed using Digest-ALL 3 (Invitrogen, Carlsbad, CA, USA) at 37°C for 20 min, twice. The sections were then sequentially dehydrated in alcohol (70%, 85%, 95% and 100%) for 2 min each and air-dried. The *KIT* probe comprised two overlapping BAC clones, C00-84L10 and RP11-586A2, labelled by random priming with digoxigenin and detected with FITC anti-digoxigenin, and co-hybridized with a spectrum orange-labelled chromosome 4 centromeric probe (CEP4; Vysis); 100 interphase nuclei were evaluated from each specimen. The cytogenetic patterns were classified as FISH-negative (no or low genomic gain: \leq four copies of *KIT* in $>40\%$ of cells), FISH-positive (high level of polysomy: \geq four copies of *KIT* in $\geq 40\%$ of cells), or gene amplification (presence of tight *KIT* gene clusters and a ratio of *KIT* and chromosome 4 cen ≥ 2 per cell, or ≥ 15 copies of *KIT* per cell in $\geq 10\%$ of analyzed cells) [16].

Results

Morphological correlates of resistance

To direct the genomic studies and to address morphological correlates for TKI resistance heterogeneity, we sampled a broad spectrum of tumour areas in each patient by evaluating several morphological parameters. Among the 57 GIST samples, the cellular morphology ranged from typical spindle cell ($n = 26$), to epithelioid ($n = 15$), to mixed ($n = 7$) to lesions with unusual ($n = 9$) features (Figure 1A–H). The spindle cell GISTs were composed of cells with pale eosinophilic fibrillary cytoplasm, ovoid nuclei and ill-defined cell borders, often with syncytial appearance (Figure 1A). GISTs with epithelioid cell morphology were composed of round cells with eosinophilic to clear cytoplasm, arranged in sheets and nests (Figure 1B). One or more samples from patients 1, 3 and 13 had a prominent perivascular, palisading (Figure 1A) or storiform growth pattern. Two tumours obtained from patient 14 showed epithelioid morphology with abrupt transformation to a pleomorphic spindle cell sarcoma (Table 1, Figure 1C and 1D). Other unusual morphologies included huge epithelioid cells with nuclear atypia (Figure 1E) and focal intracytoplasmic inclusions, as well as pleomorphic spindle cell areas. In total, nine samples with unusual morphology were collected from patients 7, 10, 12 and 14 (Table 1). Multinucleated giant cells were present in two samples obtained from patients 4 and 5 (Table 1).

Histological treatment response was heterogeneous within and between metastases selected from a given patient. Thirty-one (54.4%) samples showed minimal,

18 (31.6%) low, 7 (12.3%) moderate and only 1 (1.7%) sample showed high treatment response (Table 1). Of the samples from patients studied after progression on sunitinib, 24 (54.5%) tumours showed minimal, 15 (34.1%) low, four (9.1%) moderate and only one (2.3%) tumour showed high treatment response, respectively.

Mitotic activity was in the range 1–100 mitoses/50 HPFs. Metastases with moderate and high treatment response (eight samples) had a median mitotic activity of 4/50 HPF (range 1/50–8/50 HPFs), whereas metastases with minimal and low response (49 samples) had a median mitotic activity of 29/50 HPFs (range 1/50–100/50 HPFs). The sizes of the individual tumour nodules were in the range 0.5–35 cm (median 6.1 cm). Tumour size did not correlate with treatment effect or the frequency of detected secondary mutations (Tables 1, 2). All post-treatment samples showed robust blood vessels surrounded by smooth muscle cells/pericytes, arguing against an antiangiogenic effect of therapy. Treatment effects were noted only in the tumour parenchyma.

Heterogeneity in the immunohistochemical staining profile

KIT (CD117) staining was scored as positive in 50/57 tumour samples. All samples from patient 7 (Figure 1E) were *KIT*-negative (Figure 1F) but showed multifocal strong positivity for SMA, caldesmon (Figure 1G) and desmin (Figure 1H), whereas MYF4 staining was negative. In three patients (2, 11 and 14) a mixture of *KIT*-positive and -negative regions was observed. *KIT*-negative samples from patients 11 and 14 showed immunoreactivity with caldesmon and SMA, respectively. By contrast, the *KIT*-negative sample from patient 2 did not stain for muscle markers.

Mutational heterogeneity

Primary *KIT* mutations were detected by D-HPLC and sequencing in 11 patients (nos 1–11) (Table 1), whereas all GIST samples from the remaining three patients (nos 12–14) were *KIT* and *PDGFRA* wild-type, including four samples obtained from a patient (no. 13) with neurofibromatosis type 1 (Table 1). In all tumour samples collected from a given patient, the same primary mutation was detected. Seven patients showed a primary *KIT* mutation in exon 11 (nos 3–9) and in two patients a primary mutation in exon 9 [nos 2 and 17] or exon 13 (nos 10 and 11) was detected (Table 1).

All 57 tumour samples were screened for secondary imatinib-resistance mutations in *KIT* by D-HPLC (Table 1, Figure 2A), which has a sensitivity of $\sim 15\%$ mutant alleles. In addition, 46/57 tumour samples were screened by allele-specific PCR (AS-PCR; Table 2, Figure 2B), which has a sensitivity of 0.5% mutant alleles. By using these complimentary techniques, secondary imatinib-resistance mutations in *KIT*




Figure 1. (A) GIST with spindle cell morphology composed of cells with a pale eosinophilic fibrillary cytoplasm, ovoid nuclei and ill-defined cell borders, with syncytial appearance and palisading (patient 3). (B) GIST with epithelioid cell morphology composed of round cells with eosinophilic cytoplasm arranged in sheets (patient 2). (C) Low-power view of a GIST in the stomach wall, with abrupt transition between two morphologically different tumour components (patient 14; *KIT* and *PDGFR* wild-type). Higher-power images of this tumour are demonstrated in (D), showing epithelioid morphology (right) and pleomorphic spindle cell morphology (left). (E) GIST showing unusual morphology, with huge epithelioid cells, vesicular nuclei and prominent nucleoli (patient 7; primary *KIT* exon 11 mutation). (F) KIT-negative GIST (patient 7). (G) KIT-negative GIST showing strong cytoplasmic staining for caldesmon (patient 7). (H) KIT-negative GIST showing strong cytoplasmic staining for desmin (patient 7)

were found in 9/11 patients (82%), irrespective of the primary (exon 9, 11 or 13) *KIT* mutation. The secondary imatinib-resistant *KIT* mutations were clustered in two regions, the ATP binding pocket of the *KIT* kinase (exons 13 and 14) and in the kinase activation loop (exon 17) (Figure 2A, B). Furthermore, multiple different secondary imatinib-resistance mutations (between two and five) were found in 6/9 patients

(67%) (nos 2, 4, 5, 8, 9, 11; Tables 1, 2). In seven samples obtained from three patients (nos 2, 5, 9), two secondary imatinib-resistance *KIT* mutations were detected in one or more individual tumour samples (Tables 1, 2).

AS-PCR assays were performed for five resistance mutation hotspots [V654A, D820G, N822K (T → A and T → G) and Y823D]. In all cases

Heterogeneity of resistance mechanisms in GIST

Table 2. Summary of secondary imatinib resistant *KIT* mutations evaluated by D-HPLC and AS-PCR

Patient no.	Treatment	Location	Primary <i>KIT</i> mutations	Secondary <i>KIT</i> mutation frequencies			
				DHPLC		AS-PCR*	
				~15%		>5%	5–0.5%
1	IM	Serosal nodule (1)	Exon 9	NMF	NMF	NMF	NMF
2	IM, SU	Serosal nodule right lower quadrant	Exon 9	NMF	NMF	NMF	NMF
		Omental nodule	Exon 9	NMF	NMF	NMF	V654A
		Periumbilical tumour	Exon 9	Exon 17	N822Y	NMF	D820G
		Serosal nodule right upper quadrant (1)	Exon 9	Exon 17	N822K	N822K (T → G)	V654A
		Serosal nodule right upper quadrant (2)	Exon 9	Exon 17	N822K	N822K (T → G)	NMF
3	IM, SU	Stomach (1)	Exon 11	Exon 17	D816H	NMF	NMF
4	IM	Liver metastasis	Exon 11	Exon 18	A829P	NMF	NMF
		Stomach (1)	Exon 11	Exon 13	V654A	V654A	NMF
5	IM, SU	Liver metastasis	Exon 11	Exon 13	V654A	V654A	D820G
		Mesenteric nodule (1)	Exon 11	Exon 17	N822Y	NMF	N822K (T → A)
		Mesenteric nodule (2)	Exon 11	Exon 17	N822Y	NMF	N822K (T → G)
		Nodule attached to gallbladder	Exon 11	Exon 17	D820G	D820G	NMF
6	IM	Intra-abdominal 1	Exon 11	Exon 13	V654A	V654A	NMF
		Intra-abdominal 2	Exon 11	Exon 13	V654A	V654A	NMF
7	IM, SU	Stomach	Exon 11	NMF	NMF	NMF	NMF
		Liver metastasis	Exon 11	NMF	NMF	NMF	NMF
		Serosal nodule	Exon 11	NMF	NMF	NMF	NMF
8	IM	Abdominal tumour (1)	Exon 11	NMF	NMF	NMF	NMF
		Abdominal tumour (2)	Exon 11	Exon 17	D820Y	NMF	NMF
		Abdominal tumour (3)	Exon 11	Exon 17	D820Y	NMF	NMF
9	IM, SU	Mesenteric nodule 1	Exon 11	Exon 13	V654A	V654A	NMF
		Skin metastasis	Exon 11	Exon 13	V654A	V654A	NMF
		Mesenteric nodule 4	Exon 11	Exon 13	V654A	V654A	N822K (T → A)
10	IM	Stomach wall (3)	Exon 13	NMF	NMF	NMF	D820G
11	IM, SU	Tumour mid-abdomen	Exon 13	NMF	NMF	NMF	NMF
		Serosal nodule left upper quadrant (1)	Exon 13	Exon 14	T670I	NMF	NMF
		Serosal nodule ileum (1)	Exon 13	Exon 17	D816H	NMF	NMF
		Serosal nodule ileum (2)	Exon 13	Exon 17	D816H	NMF	NMF
		Serosal nodule terminal ileum	Exon 13	Exon 17	D816H	NMF	NMF
		Serosal nodule left upper quadrant (2)	Exon 13	Exon 17	D816H	NMF	NMF
12	IM, SU	Omental nodule					
		Liver metastasis (1)					
		Tumour sigmoid wall	NMF	NMF	NMF	NMF	
		Liver metastasis (2)					
13, NF Type I	IM, SU	Serosal nodule adjacent to bowel					
		Nodule ligamentum of Tritz					
		Nodule duodenum	NMF	NMF	NMF	NMF	
14	IM, SU	Mesenteric nodule					
		Perigastric mass (1)					
		Perigastric mass (2)					
		Perigastric mass (3)	NMF	NMF	NMF	NMF	
		Liver metastasis					
		Large nodule lesser curvature (1)					
		Large nodule lesser curvature (2)					

* AS-PCR detects: exon 13, V654A, exon 17, D820G, N822K (T → A), N822K (T → G), Y823D.
pt, patient; NMF, no mutation found.

where one of these mutations was detected by D-HPLC, there was a strong signal (>5%) by AS-PCR (Table 2). AS-PCR identified secondary *KIT* resistance mutations that were not demonstrable by D-HPLC in one or more tumour samples from four patients (nos 2, 5, 9, 10; Table 2). These additional

resistance mutations were present at low abundance (0.5–5%) in the background of more abundant (D-HPLC-detectable) secondary mutations, or as the only anomaly. Interestingly, all samples collected from patient 7 lacked *KIT* expression and neither D-HPLC nor AS-PCR revealed secondary resistance mutations




Figure 2. (A) Summary of secondary *KIT* resistance mutations detected by D-HPLC in 27/57 tumour samples from 14 patients with progressing GISTs. Associated primary *KIT* mutations are indicated in red (exon 11 mutations), yellow (exon 9 mutations) and green (exon 13 mutations). (B) Summary of secondary *KIT* resistance mutations detected by AS-PCR only (⊗) or detected by both AS-PCR and D-HPLC (●) in 18 tumour samples. Associated primary *KIT* mutations are indicated in red (exon 11 mutations), yellow (exon 9 mutations) and green (exon 13 mutations)

in these samples (Tables 1, 2). Furthermore, D-HPLC and AS-PCR did not detect secondary *KIT* resistance mutations in tumour samples showing unusual morphological features, in *KIT/PDGFR* wild-type GISTs or in *KIT*-negative GISTs (Tables 1, 2). A metastasis showing a T670I resistance mutation was the only tumour in this study with high morphological treatment response (Table 1).

Fluorescence *in situ* analysis

FISH was performed in 10/14 GIST samples showing a primary *KIT* mutation in exon 9 (nos 1, 2), exon 11 (nos 3, 7, 8) and exon 13 (nos 10, 11) but lacking a secondary *KIT* mutation. Eight of the 10 samples were FISH-negative (Figure 3A), whereas two samples, both from patient 10, had *KIT* amplification (Figure 3B).




Figure 3. (A) FISH analysis of abdominal tumour 1 (spindle cell morphology) from patient 8, with disomic (FISH-negative) pattern. The chromosome 4 centromere probe is shown in orange and the *KIT* probe in green. (B) FISH analysis of stomach wall (2) tumour (epithelioid morphology) from patient 10, with *KIT* amplification. Chromosome 4 centromere probe is shown in orange and the *KIT* probe in green

Discussion

The aim of this study was to determine the heterogeneity of tyrosine kinase inhibitor (TKI) resistance mutations within and between different metastatic GIST lesions and to examine the relationship between drug resistance and morphological variability, mitotic activity and immunophenotype. Secondary *KIT* mutations were interrogated using two complementary techniques, D-HPLC and AS-PCR. D-HPLC provided a broad unbiased screening approach to detect secondary mutations in various *KIT* and *PDGFRA* exons, with a sensitivity of ~15%. By contrast, the novel AS-PCR approach was designed to detect five hotspot secondary imatinib-resistant *KIT* mutations, with a sensitivity of ~0.5%. Secondary *KIT* kinase domain mutations were found in 82% of patients after imatinib or sunitinib therapy, irrespective of whether the primary

Heterogeneity of resistance mechanisms in GIST

mutation was in *KIT* exon 9, 11 or 13. The secondary mutations were clustered in the ATP binding pocket (D-HPLC 42.9% and AS-PCR 50%) and in the activation loop (D-HPLC 57.1% and AS-PCR 50%) of the *KIT* kinase domain. D-HPLC demonstrated two alternative secondary *KIT* resistance mutations (involving exons 13 and 17) in only 1/56 tumour samples, whereas the more sensitive AS-PCR method demonstrated more than one resistance mutation in each of six metastases. These findings underscore the complexity of clinically important TKI resistance mechanisms. Whereas previous reports have demonstrated resistance mutations in 44–67% of GISTs progressing after imatinib therapy [6,17–19], the combination of D-HPLC and AS-PCR revealed secondary *KIT* TKI-resistance mutations in 9/10 patients (90%) whose GIST had a primary *KIT* oncogenic mutation and whose tumour expressed *KIT* protein. In addition, we demonstrated two or more (between two and five) TKI-resistance mutations in most patients in this series, whereas previous studies have shown multiple resistance mutations in ~10–30% of GIST patients at the time of clinical progression [6,19]. Contributing factors to the higher level of demonstrable TKI-resistance mutations might include the more extensive sampling of multiple metastases, incorporation of the highly sensitive AS-PCR detection method, and clonal selection for additional resistance mutations in patients receiving sunitinib. Patients receiving sunitinib treatment showed substantially more (one to five) secondary imatinib-resistance mutations compared to patients only treated with imatinib (at most two). However, the mutation types seen after sunitinib therapy, in this study, were similar to those reported after progression on imatinib therapy alone [6,8,10]. Secondary resistance mutations in the *KIT* activation loop (exon 17) were seen in ~60% (Table 2) of sunitinib-treated metastases, which is in accord with *in vitro* evidence that mutations in this region are sunitinib-resistant [20]. Although multiple *KIT* exons (9,11–18) were screened for secondary mutations outside the ATP binding pocket and the kinase activation loop, D-HPLC did not reveal mutations in these regions.

KIT V654A is the most frequent secondary mutation in patients whose GISTs have primary *KIT* exon 11 mutations and who eventually progress during imatinib treatment [6,8]. Interestingly, the V654A mutation, which is sunitinib-sensitive based on *in vitro* studies [20], was found in ~27% (Table 2) of samples after clinical progression on sunitinib. In addition, these same samples showed minimal to low morphological evidence of treatment response. Such observations suggest that sunitinib might be cytostatic rather than cytotoxic in GISTs with secondary V654A mutations. This is in keeping with the clinical evidence from a randomized, placebo-controlled, multicentre trial demonstrating that stable GIST was the best overall tumour response on sunitinib treatment [11]. In addition, the presence of low-level TKI resistance mutations on the same *KIT* alleles encoding

the V654A might also account for the persistence of V654A alleles during sunitinib therapy.

In theory, sunitinib therapeutic activity could result from either antiangiogenic or direct antitumoral activity, as this is a multikinase inhibitor that inhibits *KIT*, *PDGFR*, *VEGFR*, *FLT3* and *RET* [21–25]. However, there is no compelling evidence for sunitinib antiangiogenic activity in GIST patients [26]. In the present study, moderate-to-high morphological evidence of sunitinib treatment response was found in 11.4% of metastases, even at time of clinical progression on sunitinib. Viable tumour cells from these samples were often clustered around blood vessels, and a reduction of overall tumour vasculature was not detected, suggesting that the treatment response in these cases resulted largely from direct inhibition of crucial *KIT*-mediated survival pathways in the GIST cells, rather than anti-angiogenic effects. As expected, GIST metastases with moderate and high treatment response had lower mitotic activity overall (median 4/50HPF) compared to those with minimal or low treatment response (median 29/50HPF). However, there was substantial variability in the mitotic activity amongst samples with minimal and low treatment response, such that mitotic activity alone is not a reliable marker for clinical progression. Interruption of TKI therapy, as might occur pre-operatively, is known to augment *KIT* phosphorylation and the biochemical activity of downstream signalling proteins in GIST, and to cause PET scan 'flares' [17]. Therefore, evaluation of treatment effects based on more fixed morphological correlates, such as necrosis and fibrosis, might be a more reliable indicator than mitotic activity in distinguishing clinically stable from progressing GIST lesions. In our study, tumour size did not correlate with the presence of treatment effect, and this has been noted in a previous report [27]. Furthermore, there was no correlation between tumour size and frequency of detectable secondary *KIT* kinase mutations.

Using complementary D-HPLC and AS-PCR mutational screening approaches, we demonstrated secondary resistance mutations as the overriding resistance mechanism in 9/14 patients. By contrast, all patients whose GISTs had wild-type *KIT/PDGFR* lacked kinase domain resistance mutations, indicating that mechanisms of resistance are fundamentally different in these GISTs. *KIT* wild-type GISTs feature levels of *KIT* activation similar to *KIT*-mutant GISTs [28,29], suggesting that *KIT* has a key oncogenic role in the pathogenesis of these tumours. However, imatinib is a less potent inhibitor of wild-type *KIT* compared with exon 11-mutant *KIT* [34, Heinrich *et al*, in press], possibly accounting for the lower clinical imatinib responsiveness for GISTs expressing wild-type *KIT*.

Unusual morphologies were seen in four patients, including two whose GISTs were *KIT/PDGFR* wild-type and two with metastases that lacked *KIT* expression. The distinctive morphological features included

large epithelioid cells with vesicular chromatin, prominent nucleoli and focal intracytoplasmic eosinophilic inclusions, and abrupt transformation/dedifferentiation in a high-grade spindle cell tumour. Mutational analyses of samples entirely comprised of these unusual morphological features did not reveal novel *KIT* mutations as the potential molecular mechanism accounting for these bizarre histological evolutions. However, the observed loss of KIT expression in some of these samples suggests that the histological changes resulted from activation of novel KIT-independent oncogenic pathways. Interestingly, smooth muscle markers were expressed strongly in six of the seven samples featuring loss of KIT expression. These observations fit with developmental biology evidence that smooth muscle cells and interstitial cells of Cajal (ICC) arise from a common progenitor cell, with KIT signalling committing these progenitor cells to ICC differentiation, and KIT inhibition resulting in a switch to smooth muscle differentiation [30]. Therefore, unusual morphological and immunohistochemical features are found in GISTs after TKI treatment, and might present diagnostic challenges, particularly if coupled with loss of KIT expression, as detected by IHC.

Our studies also confirm a role, albeit limited [31], for *KIT* amplification as a mechanism of drug resistance in GIST. Analyses in 10 GISTs lacking demonstrable secondary *KIT* mutations revealed localized *KIT* amplicons in both samples analysed from patient 10, whose GIST had a primary *KIT* exon 13 mutation. These findings suggest that increased gene dosage can contribute to clinical progression in some GISTs.

Overall, the molecular, FISH and histological assessments described here underscore a striking inter- and intra-lesional heterogeneity in TKI resistance mechanisms. With regard to treatment approaches, newer generations of broad-spectrum KIT and PDGFRA kinase inhibitors, as well as combination therapies with various such inhibitors, could prolong GIST remissions in a manner similar to treatment approaches used in HIV, by suppressing a broader spectrum of tumour clones from the outset of therapy. Novel treatment options include inhibition of the KIT chaperone HSP90 [32], which may result in KIT oncoprotein degradation, irrespective of the TKI-resistance mutations present. Other broadly relevant therapeutic strategies include blockage of crucial KIT-mediated signalling pathways, as might be accomplished via PI3-K inhibition [33].

In summary, our study underscores the overriding contribution of secondary *KIT* mutations to TKI resistance in GISTs and demonstrates substantial heterogeneity of resistance mutations within and between metastases from individual patients. However, the highly sensitive and complementary D-HPLC and

AS-PCR methods did not show TKI-resistance mutations in tumours wild-type for *KIT/PDGFR*A, indicating that resistance mechanisms are fundamentally different in these GISTs. Likewise, secondary *KIT* TKI-resistance mutations were not found in clinically progressing GISTs that lacked KIT expression, suggesting that activation of novel oncogenic pathways accounts for TKI resistance in such cases. Our observations on the complexity of TKI resistance underscore the challenges in achieving long-term disease control with kinase inhibitor monotherapies, and raise concern over the ultimate effectiveness of second- and third-line TKI drugs in GIST patients resistant to imatinib.

Acknowledgements

This work was supported by grants from an anonymous donor, GI SPORE 1P50CA127003-01, the Life Raft Group, the Cesarini Team for the Pan-Massachusetts Challenge, the Virginia and Daniel K. Ludwig Trust for Cancer Research, the Ronald O. Perelman Fund for Cancer Research, the Stutman GIST Cancer Research Fund, the BP Lester Foundation, the Rubenstein Foundation, Leslie's Links, a VA Merit Review Grant and the FWF Austrian Science Fund.

References

1. Heinrich MC, Corless CL, Demetri GD, Blanke CD, von Mehren M, Joensuu H, *et al*. Kinase mutations and imatinib response in patients with metastatic gastrointestinal stromal tumor. *J Clin Oncol* 2003;**21**(23):4342–4349.
2. Heinrich MC, Corless CL, Duensing A, McGreevey L, Chen CJ, Joseph N, *et al*. PDGFRA activating mutations in gastrointestinal stromal tumors. *Science* 2003;**299**(5607):708–710.
3. Hirota S, Isozaki K, Moriyama Y, Hashimoto K, Nishida T, Ishiguro S, *et al*. Gain-of-function mutations of *c-kit* in human gastrointestinal stromal tumors. *Science* 1998;**279**(5350):577–580.
4. Demetri GD, von Mehren M, Blanke CD, Van den Abbeele AD, Eisenberg B, Roberts PJ, *et al*. Efficacy and safety of imatinib mesylate in advanced gastrointestinal stromal tumors. *N Engl J Med* 2002;**347**(7):472–480.
5. Verweij J, Casali PG, Zalcberg J, LeCesne A, Reichardt P, Blay JY, *et al*. Progression-free survival in gastrointestinal stromal tumours with high-dose imatinib: randomised trial. *Lancet* 2004;**364**(9440):1127–1134.
6. Heinrich M, Corless C, Blanke C, *et al*. Molecular correlates of imatinib resistance in gastrointestinal stromal tumors. *J Clin Oncol* 2006;**24**(29):4764–4774.
7. Debiec-Rychter M, Dumez H, Judson I, Wasag B, Verweij J, Brown M, *et al*. Use of *c-KIT/PDGFR*A mutational analysis to predict the clinical response to imatinib in patients with advanced gastrointestinal stromal tumours entered on phase I and II studies of the EORTC Soft Tissue and Bone Sarcoma Group. *Eur J Cancer* 2004;**40**(5):689–695.
8. Antonescu CR, Besmer P, Guo T, Arkun K, Hom G, Korytowski B, *et al*. Acquired resistance to imatinib in gastrointestinal stromal tumor occurs through secondary gene mutation. *Clin Cancer Res* 2005;**11**(11):4182–4190.
9. Fletcher J, Corless C, Dimitrijevic S, *et al*. Mechanisms of resistance to imatinib mesylate (IM) in advanced gastrointestinal stromal tumors (GIST). *Proc Am Soc Clin Oncol* 2003;**22**:3275–3277.
10. Debiec-Rychter M, Cools J, Dumez H, Sciot R, Stul M, Mentens N, *et al*. Mechanisms of resistance to imatinib mesylate in gastrointestinal stromal tumors and activity of the PKC412 inhibitor against imatinib-resistant mutants. *Gastroenterology* 2005;**128**(2):270–279.

Heterogeneity of resistance mechanisms in GIST

- Demetri GD, van Oosterom AT, Garrett CR, Blackstein ME, Shah MH, Verweij J, *et al.* Efficacy and safety of sunitinib in patients with advanced gastrointestinal stromal tumour after failure of imatinib: a randomised controlled trial. *Lancet* 2006;**368**(9544):1329–1338.
- Therasse P, Le Cesne A, Van Glabbeke M, Verweij J, Judson I. RECIST vs. WHO: prospective comparison of response criteria in an EORTC phase II clinical trial investigating ET-743 in advanced soft tissue sarcoma. *Eur J Cancer* 2005;**41**(10):1426–1430.
- Hornick JL, Fletcher CD. Immunohistochemical staining for KIT (CD117) in soft tissue sarcomas is very limited in distribution. *Am J Clin Pathol* 2002;**117**(2):188–193.
- Corless CL, McGreevey L, Haley A, Town A, Heinrich MC. KIT mutations are common in incidental gastrointestinal stromal tumors one centimeter or less in size. *Am J Pathol* 2002;**160**(5):1567–1572.
- Corless CL, Schroeder A, Griffith D, Town A, McGreevey L, Harrell P, *et al.* PDGFRA mutations in gastrointestinal stromal tumors: frequency, spectrum and *in vitro* sensitivity to imatinib. *J Clin Oncol* 2005;**23**(23):5357–5364.
- Cappuzzo F, Varella-Garcia M, Shigematsu H, Domenichini I, Bartolini S, Ceresoli GI, *et al.* Increased HER2 gene copy number is associated with response to gefitinib therapy in epidermal growth factor receptor-positive non-small-cell lung cancer patients. *J Clin Oncol* 2005;**23**(22):5007–5018.
- Agaram NP, Besmer P, Wong GC, Guo T, Socci ND, Maki RG, *et al.* Pathologic and molecular heterogeneity in imatinib-stable or imatinib-responsive gastrointestinal stromal tumors. *Clin Cancer Res* 2007;**13**(1):170–181.
- Prenen H, Cools J, Mentens N, Folens C, Sciot R, Schoffski P, *et al.* Efficacy of the kinase inhibitor SU11248 against gastrointestinal stromal tumor mutants refractory to imatinib mesylate. *Clin Cancer Res* 2006;**12**(8):2622–2627.
- Wardelmann E, Merkelbach-Bruse S, Pauls K, *et al.* Polyclonal evolution of multiple secondary KIT mutations in gastrointestinal stromal tumors under treatment with imatinib mesylate. *Clin Cancer Res* 2006;**12**(6):1743–1749.
- Heinrich MC, Corless CL, Liegl B, Fletcher CD, Raut CP, Donsky R, *et al.* Mechanisms of sunitinib malate (SU) resistance in gastrointestinal stromal tumors (GISTs). *J Clin Oncol* 2007;**25**(1; No. 18S, June 20 suppl): Abstract No: 10006.
- Osusky KL, Hallahan DE, Fu A, Ye F, Shyr Y, Geng L. The receptor tyrosine kinase inhibitor SU11248 impedes endothelial cell migration, tubule formation, and blood vessel formation *in vivo*, but has little effect on existing tumor vessels. *Angiogenesis* 2004;**7**(3):225–233.
- Abrams TJ, Lee LB, Murray LJ, Pryer NK, Cherrington JM. SU11248 inhibits KIT and platelet-derived growth factor receptor beta in preclinical models of human small cell lung cancer. *Mol Cancer Ther* 2003;**2**(5):471–478.
- Mendel DB, Laird AD, Xin X, Louie SG, Christensen JG, Li G, *et al.* *In vivo* antitumor activity of SU11248, a novel tyrosine kinase inhibitor targeting vascular endothelial growth factor and platelet-derived growth factor receptors: determination of a pharmacokinetic/pharmacodynamic relationship. *Clin Cancer Res* 2003;**9**(1):327–337.
- O'Farrell AM, Abrams TJ, Yuen HA, Ngai TJ, Louie SG, Yee KW, *et al.* SU11248 is a novel FLT3 tyrosine kinase inhibitor with potent activity *in vitro* and *in vivo*. *Blood* 2003;**101**(9):3597–3605.
- Schueneman AJ, Himmelfarb E, Geng L, Tan J, Donnelly E, Mendel D, *et al.* SU11248 maintenance therapy prevents tumor regrowth after fractionated irradiation of murine tumor models. *Cancer Res* 2003;**63**(14):4009–4016.
- Seandel M, Shia J, Linkov I, Maki RG, Antonescu CR, Dupont J. The activity of sunitinib against gastrointestinal stromal tumor seems to be distinct from its antiangiogenic effects. *Clin Cancer Res* 2006;**12**(20):part 1: 6203–6204.
- Agaram NP, Baren A, Arkun K, Dematteo RP, Besmer P, Antonescu CR. Comparative ultrastructural analysis and KIT/PDGFR α genotype in 125 gastrointestinal stromal tumors. *Ultrastruct Pathol* 2006;**30**(6):443–452.
- Rubin BP, Singer S, Tsao C, Duensing A, Lux ML, Ruiz R, *et al.* KIT activation is a ubiquitous feature of gastrointestinal stromal tumors. *Cancer Res* 2001;**61**(22):8118–8121.
- Janeway KA, Liegl B, Harlow A, Le C, Perez-Atayde A, Kozakewich H, *et al.* Pediatric KIT wild-type and platelet-derived growth factor receptor alpha-wild-type gastrointestinal stromal tumors share KIT activation but not mechanisms of genetic progression with adult gastrointestinal stromal tumors. *Cancer Res* 2007;**67**(19):9084–9088.
- Torihashi S, Nishi K, Tokutomi Y. Blockade of KIT signaling induces transdifferentiation in interstitial cells of Cajal to smooth muscle phenotype. *Gastroenterology* 1999;**117**:140–148.
- Miselli F, Casieri P, Negri T, Orsenigo M, Lagonigro MS, Gronchi A, *et al.* c-KIT/PDGFR α gene status alterations possibly related to primary imatinib resistance in gastrointestinal stromal tumors. *Clin Cancer Res* 2007;**13**(8):2369–2377.
- Bauer S, Yu LK, Demetri GD, Fletcher JA. Heat shock protein 90 inhibition in imatinib-resistant gastrointestinal stromal tumor. *Cancer Res* 2006;**66**(18):9153–9161.
- Bauer S, Duensing A, Demetri GD, Fletcher JA. KIT oncogenic signaling mechanisms in imatinib-resistant gastrointestinal stromal tumor: PI3-kinase/AKT is a crucial survival pathway. *Oncogene* 2007;**26**(54):7560–7568.
- Heinrich MC, Maki RG, Corless CL, Antonescu CR, Horlow A, Griffith D, *et al.* Primary and secondary kinase genotypes correlate with the biological and clinical activity of sunitinib malate in imatinib-resistant gastrointestinal stromal tumours. *J Clin Oncol*; in press.